

**Problem Set 6.**

Bosons and fermions.

**Problem 6.1** Coherent states of bosons.

For the coherent state of bosons

$$|a\rangle = e^{-\frac{|a|^2}{2}} \sum_{n=0}^{\infty} \frac{a^n}{\sqrt{n!}} |n\rangle \quad (1)$$

(1) Prove that

$$\langle a|d|a\rangle = \frac{1}{2}(a^* da - a da^*) \quad (2)$$

( $d$  denotes differentiation)

(2) For a closed contour  $\Gamma$  in the complex plane ( $a, a^*$ ), prove that the integral

$$\int_{\Gamma} \langle a|d|a\rangle \quad (3)$$

is proportional to the area inside  $\Gamma$ .

**Problem 6.2** Consider the ground state of one-dimensional spinless non-interacting fermions. The density of fermions is  $n$ .

(1) Find the equal-time Green's function in the ground state

$$\langle c^\dagger(x)c(y)\rangle \quad (4)$$

( $c^\dagger$  and  $c$  are the creation and annihilation operators for the fermions). Note that it does not depend on the particle mass.

(2) Find the equal-time density-density correlation in the ground state

$$\langle \rho(x)\rho(y)\rangle, \quad \text{where } \rho(x) = c^\dagger(x)c(x) \quad (5)$$

(3) Consider a large interval of length  $L$ . Define the number of particles in this interval as

$$N = \int_0^L \rho(x) dx \quad (6)$$

On average, the particle number in this interval is  $\langle N\rangle = nL$ .

Calculate the quantum fluctuations of  $N$  in the ground state

$$(\delta N)^2 = \langle N^2\rangle - \langle N\rangle^2 \quad (7)$$

Show that at large  $N$  the fluctuation  $(\delta N)^2$  grows as  $\ln N$ . This contrasts the situation with the gas of classical particles, where the distribution of  $N$  is Poissonian with  $(\delta N)^2 \sim N$ .